

MAR10-2009-006797

Abstract Submitted  
for the MAR10 Meeting of  
The American Physical Society

Sorting Category: 14.7 (C)

**Molecular Dynamics**  
**simulations of tetracosane ( $C_{24}H_{50}$ ) bilayers physisorbed onto the basal plane of graphite**<sup>1</sup> MICHAEL ROTH, E. MALDONADO, University of Northern Iowa, L. FIRLEJ, LCVN, Universit Montpellier 2, B. KUČHTA, Laboratoire Chimie Provence, Universit de Provence, CARLOS WEXLER, University of Missouri Department of Physics and Astronomy — We present and discuss the results of explicit - hydrogen Molecular Dynamics computer simulations of tetracosane ( $C_{24}H_{50}$ ) bilayers deposited on a graphite substrate in the temperature range  $100\text{ K} \leq T \leq 450\text{ K}$ . Both layers exhibit strong coupling between the internal molecular degrees of freedom and bulk behavior but because of the different boundary conditions between layers, they exhibit distinctly different dynamics and phase transition signatures. Structural, thermodynamic and bond - orientational distributions and parameters are utilized in understanding the solid, intermediate and liquid phases presented in and phase transitions presented by the system.

<sup>1</sup>Donors of The American Chemical Society Petroleum Research Fund (PRF43277-B5), University of Missouri Research Board, Department of Energy award number DE-FG02-07ER46411

Prefer Oral Session  
 Prefer Poster Session

Michael Roth  
rothm@uni.edu  
University of Northern Iowa

Date submitted: 20 Nov 2009

Electronic form version 1.4