

MAR09-2008-005241

Abstract Submitted
for the MAR09 Meeting of
The American Physical Society

Sorting Category: 04.11 (C)

**All-atom Molecular Dynamics simulations of partial pentane
and hexane films on graphite**

¹ M KASPAR, M.W. ROTH, University of Northern Iowa, CARLOS WEXLER, University of Missouri, L. FIRLEJ, Universit Montpellier, B. KUCHTA, Universit de Provence — We compare the self-assembly patterns of pentane (C_5H_{12}) and hexane (C_6H_{14}) adlayers physisorbed onto graphite at various coverages using the results of molecular dynamics simulations. Near monolayer coverage, the solid low temperature structure of the pentane film is nematic-like, and that of hexane-herringbone-like. At submonolayer coverages both systems exhibit three distinct topological regimes: vacancy patches at higher densities, percolating networks at intermediate densities and ultimately individual patches. The systems' orientational behavior and melting dynamics is discussed with respect to its unique density-dependent topology. The simulations explicitly include hydrogens of pentane and hexane and the graphite is modeled as a six-layer all atom structure.

¹Work supported by the U.S. Department of Energy (DE-FG02-07ER46411) and the American Chemical Society Petroleum Research Fund (PRF43277-B5). Computational resources were provided by the University of Missouri Bioinformatics Consortium.

Prefer Oral Session
 Prefer Poster Session

Michael Roth
rothm@uni.edu
University of Northern Iowa

Date submitted: 21 Nov 2008

Electronic form version 1.4