

MAR08-2007-007146

Abstract Submitted
for the MAR08 Meeting of
The American Physical Society

Sorting Category: 14.7 (C)

**Test of the universality of the scaling energy in alkanes using
melting transition of layer adsorbed on graphite**

¹ B. KUCHTA, L. FIRLEJ, M.W. ROTH, ¹University of Northern Iowa, Department of Physics, PAUL A. GRAY, University of Northern Iowa, Department of Computer Science, CARLOS WEXLER, ³University of Missouri Columbia, Department of Physics and Astronomy — The non-bonded terms together with the torsional energy determine the internal (conformational) degrees of freedom of simulated alkanes. Being able to predict the energy and ordering of conformations correctly is an essential quantity for bringing force field methods into the predictive regime of theoretical methods. To estimate the universality of modern force fields it is helpful to know how different components of classical fields affect the simulated properties. In the case of alkanes, the interplay between torsion potential and the scaling of 1-4 van der Waals and 1-4 charge-charge (QQ) interactions plays a crucial role. It is the purpose of this work to study universality of the scaling factor of 1-4 non-bonded interactions in alkanes. Three alkanes of length from 7 to 30Å (C₆, C₁₂, C₂₄), in an all-atom representation have been analyzed.

¹Acknowledgment is made to the Donors of The American Chemical Society Petroleum Research Fund (PRF43277 B5), University of Missouri Research Board, and the D.O.E. (DE-FG02-07ER46411).

Prefer Oral Session
 Prefer Poster Session

Michael Roth
rothm@uni.edu
University of Northern Iowa

Date submitted: 27 Nov 2007

Electronic form version 1.4